

ASSIGNMENT – 1  
M.Sc. DEGREE EXAMINATION, JUNE/JULY - 2020  
(Second Year)  
MATHEMATICS  
Topology and Functional Analysis  
MAXIMUM MARKS – 30  
ANSWER ALL QUESTIONS

- Q1)** a) Show that a subspace of a topological space is itself a topological space.  
b) Show that a topological space  $X$  is metrizable there exist a homeomorphism of  $X$  onto a subspace of some metric space  $Y$ .
- Q2)** a) State and prove Tychonoff's Theorem.  
b) Prove that every sequentially compact metric space is totally bounded.
- Q3)** a) State and Prove Ascoli's theorem.  
b) Show that  $\mathbb{R}^{\mathbb{R}}$  is not locally compact.
- Q4)** a) Prove that the product of any non-empty class of Hausdorff space is a Hausdorff space.  
b) Prove that compact subspace of a Hausdorff space is closed.
- Q5)** a) State and prove the Urysohn Imbedding theorem.  
b) State and prove a generalization of Tietze's theorem which relates to functions whose values lie in  $\mathbb{R}^n$

**ASSIGNMENT – 2**  
**M.Sc. DEGREE EXAMINATION, JUNE/JULY - 2020**  
**(Second Year)**  
**MATHEMATICS**  
**Topology and Functional Analysis**  
**MAXIMUM MARKS – 30**  
**ANSWER ALL QUESTIONS**

**Q1)**

- a) Define Banach Space and give some examples.
- b) Let  $N$  be a non-zero normed linear space, and prove that  $N$  is a Banach space  $\Leftrightarrow \{x: \|x\|=1\}$  is complete.

- a) State and prove the Hahn-Banach Theorem.
- b) If  $N$  is a normed linear space and  $x_0$  is a non-zero vector in  $N$ , then there exist a functional  $f_0$  in  $N^*$  such that  $f_0(x_0) = \|x_0\|$  and  $\|f_0\|=1$ .

**Q2)**

- Q3)** a) Prove that a closed convex subset  $C$  of a Hilbert space  $H$  contains a unique vector of smallest norm.
- b) Define Hilbert Space and give some examples.

**Q4)** a) State and prove Bessel's inequality.

- b) Prove that a Hilbert space  $H$  is separable if and only if every orthonormal set in  $H$  is countable.

**Q5)** a) If  $T$  is an operator on  $H$  for which  $(T_{x,x}) = 0$  for all  $x$ , then prove that  $T = 0$ .

- b) If  $T$  is an operator on  $H$ , then prove that the following conditions are all equivalent to each other.

- i)  $TT^* = I$ .
- ii)  $(T_x, T_y) = (x, y)$  for all  $x$  and  $y$ .
- iii)  $\|Tx\| = \|x\|$  for all  $x$ .

**ASSIGNMENT – 1**  
**M.Sc. DEGREE EXAMINATION, JUNE/JULY - 2020**  
**(Second Year)**  
**MATHEMATICS**  
**Measure and Integration**  
**MAXIMUM MARKS – 30**  
**ANSWER ALL QUESTIONS.**

- Q1)** a) The union of a countable collection of countable sets is countable.  
 b) Show that every bounded infinite sequence has a subsequence that converges to a real number.
- Q2)** a) Let  $\{A_n\}$  be a countable collection of sets of real numbers, then prove that  $m^*(\cup A_n) \leq \sum m^* A_n$ .  
 b) If  $f$  is measurable function and  $f = g$  a.e, then prove that  $g$  is measurable.
- Q3)** a) State and prove Egoroff's theorem.  
 b) State and prove Lusin's theorem.
- Q4)** a) Let  $\phi$  and  $\psi$  be simple functions which vanish outside a set of finite measure, then prove that  $\int (a\phi + b\psi) = a\int \phi + b\int \psi$ .  
 b) State and prove Fatou's Lemma.
- Q5)** a) Show that if  $f$  is integrable over  $E$ , then so is  $|f|$  and  $|\int_E f| \leq \int_E |f|$   
 b) Let  $\langle f_n \rangle$  be a sequence of measurable functions that converges in measure to  $f$ . Then prove that, there is subsequence  $\langle f_{n_k} \rangle$  that converges to  $f$  almost everywhere.

**ASSIGNMENT – 2**  
**M.Sc. DEGREE EXAMINATION, JUNE/JULY - 2020**  
**(Second Year)**  
**MATHEMATICS**  
**Measure and Integration**  
**MAXIMUM MARKS – 30**  
**ANSWER ALL QUESTIONS.**

Q1)

a) Let  $f$  be an increasing real-valued function on the interval  $[a, b]$ . Then prove that  $f$  is differentiable almost everywhere. The derivative  $f'$  is measurable, and  $\int_a^b f'(x) dx \leq f(b) - f(a)$ .

b) If  $f$  is integrable on  $[a, b]$ , then show that the function  $F$  defined by  $F(x) = \int_a^x f(t) dt$  is a continuous function of bounded variation on  $[a, b]$ .

Q2)

- a) Prove the Minkowski Inequality for  $1 \leq p < \infty$ .
- b) Given  $f \in L^p$ ,  $1 \leq p < \infty$ , and  $\varepsilon > 0$ , then prove that there is a bounded measurable function  $f_M$  with  $|f_M| \leq M$  and  $\|f - f_M\| < \varepsilon$ .

Q3)

- a) If  $E_i \in \mathfrak{E}$ ,  $\mu E_1 < \infty$  and  $E_i \supset E_1$ , then prove that 
$$\mu \left( \bigcap_{i=1}^{\infty} E_i \right) = \lim_{n \rightarrow \infty} \mu E_n.$$
- b) Suppose that to each  $\alpha$  in a dense set  $D$  of real numbers there is assigned a set  $B_\alpha \in \mathfrak{E}$  such that  $B_\alpha \subset B_\beta$  for  $\alpha < \beta$ . Then prove that there is unique measurable extended real valued function  $f$  on  $X$  such that  $f \leq \alpha$  on  $B_\alpha$  and  $f \geq \alpha$  on  $X \sim B_\alpha$ .

Q4) State and prove Hahn Decomposition theorem.

Q5)

The class of  $\mathfrak{E}$  of  $\mu^*$  - measurable sets is a  $\sigma$ -algebra.

If  $\bar{\mu}$  is  $\mu^*$  restricted to  $\mathfrak{E}$ , then prove that  $\bar{\mu}$  is complete measure on  $\mathfrak{E}$ .



ASSIGNMENT – 1  
M.Sc. DEGREE EXAMINATION, JUNE/JULY - 2020

(Second Year)

MATHEMATICS

Analytical Number Theory and Graph Theory

MAXIMUM MARKS – 30

ANSWER ALL QUESTIONS

**Q1)**

a) For all  $x \geq 1$ , Prove that  $\sum_{n \leq x} \sigma_1(n) = \frac{1}{2} \zeta(2) x^2 + O(x \log x)$ . Prove that  $\sum_{n \leq x} \sigma_\alpha(n) = \frac{\zeta(\alpha+1)}{\alpha+1} x^{\alpha+1} + O(x^\alpha)$ , where  $\beta = \max\{1, \alpha\}$

b) State and prove Euler's summation formula.

**Q2)**

a) For all  $x \geq 1$ , prove that  $\left| \sum_{n \leq x} \frac{\mu(n)}{n} \right| \leq 1$  with equality holding only if  $x < 2$ .

b) State and prove Legendre's identity.

**Q3)** a) State and prove Abel's identity.

b) For a  $x \geq 2$ , prove that  $v(x) = \pi(x) \log x - \int_2^x \frac{\pi(t)}{t} dt$

$$\text{and } \pi(x) = \frac{v(x)}{\log x} + \int_2^x \frac{v(t)}{t \log^2 t} dt.$$

a) Prove that the prime number theorem implies

$$\lim_{x \rightarrow \infty} \frac{M(x)}{x} = 0.$$

**Q4)**

b) State and prove Selberg's asymptotic formula.

**Q5)** a) Prove that, a simple with  $n$  vertices and  $k$  components can have at most  $(n - k)(n - k + 1)/2$  edges.

b) Prove that, a connected graph  $G$  is an Euler graph if and only if it can be decomposed into circuits.

ASSIGNMENT – 2  
M.Sc. DEGREE EXAMINATION, JUNE/JULY - 2020

(Second Year)

MATHEMATICS

Analytical Number Theory and Graph Theory

MAXIMUM MARKS – 30

ANSWER ALL QUESTIONS

- Q1)** a) Explain Konigsberg Bridge problem.  
b) Prove that, an Euler graph  $G$  is arbitrary traceable from vertex  $v$  in  $G$  if and only if every circuit in  $G$  contains  $v$ .
- Q2)** a) If in a graph  $G$  there is one and only one path between every pair of vertices, then  $G$  is a tree.  
b) Prove that the distance between vertices of a connected graph is a metric.
- Q3)** a) Prove that the ring sum of any two cut-sets in a graph is either a third cut-set or an edge-disjoint union of cut-sets.  
b) Prove that every cut-set in a connected graph  $G$  must contain at least one branch of every spanning tree of  $G$ .
- Q4)** a) Prove that Kuratowski's second graph is non-planar.  
b) Prove that, a connected graph with  $n$  vertices and  $e$  edges has  $e - n + 2$  regions.
- Q5)** a) Prove that, the ring sum of two circuits in a graph  $G$  is either a circuit or an edge-disjoint union of circuits.  
b) Explain about Modular arithmetic and Galois fields.



**ASSIGNMENT – 1**  
**M.Sc. DEGREE EXAMINATION, JUNE/JULY - 2020**  
**(Second Year)**  
**MATHEMATICS**  
**Rings and Modules**  
**MAXIMUM MARKS – 30**  
**ANSWER ALL QUESTIONS**

**Q1)**

- a) The class of semi-lattices can be equationally defined as the class of all semi-groups  $(S, \wedge)$ , then prove that  $(S, \wedge)$  satisfying the commutative and idempotent laws.
- b) Prove that a Boolean algebra  $(S, 0', \cdot)$  can be turned into a Boolean ring  $(S, 0, 1, -, +, \cdot)$  by defining  $1 = 0'$ ,  $-a = a$ ,  $a + b = ab' \vee ba'$  where  $a \vee b = (a' b')'$ .

- a) Prove that the sum  $\sum_{i \in I} B_i$  of submodules of  $A_R$  is direct if and only if, for all  $i \in I$ ,  $B_i \cap \sum_{j \neq i} B_j = 0$ .

**Q2)**

- b) Prove that the following statements are equivalent.
- (1)  $R$  is isomorphic to a finite direct of rings  $R_i (i = 1, 2, \dots, n)$
  - (2) There exist central orthogonal idempotents,  $e_i \in R$  such that  $1 = \sum_{i=1}^n e_i$  and  $e_i R \cong R_i$ .
  - (3)  $R$  is a finite direct sum of ideals  $K_i \cong R_i$ .

**Q3)**

- a) If  $B$  and  $C$  are sub-modules of  $A$  prove that  $(B+C)/B \cong C/(B \cap C)$ .
- b) Prove that a module is Noetherian if and only if every sub-module is finitely generated.

**Q4)** a) Prove that radical of a commutative ring  $R$  consists of all nilpotent elements of.

- b) Prove the following statements concerning the Boolean ideal  $K$  are equivalent.
- (1)  $K$  is maximal
  - (2)  $K$  is prime



(3) For every element  $s$ , either  $s \in K$  or  $s' \in K$   
but not both.

- Q5)** a) Prove that every equivalent class of fractions contains exactly one irreducible fraction, and this extends all fractions in the class.
- b) If  $R$  is a Boolean ring then prove that  $Q(R)$  is a Boolean ring.

**ASSIGNMENT – 2**  
**M.Sc. DEGREE EXAMINATION, JUNE/JULY - 2020**  
**(Second Year)**  
**MATHEMATICS**  
**Rings and Modules**  
**MAXIMUM MARKS – 30**  
**ANSWER ALL QUESTIONS**

- Q1)** a) Prove that, the ring  $R$  is prime if and only if there exist a faithful irreducible module  $A_R$ .
- b) Show that a dense sub-ring of the ring of linear transformations of a vector space is primitive.
- Q2)** a) Show that, the prime radical of  $R$  is the set of all strongly nilpotent elements.
- b) Prove that, the radical of  $R$  is the set of all  $r \in R$  such that  $1 - rs$  is right invertible for all  $s \in R$ .
- Q3)** a) Prove the following conditions concerning the module  $A$  are equivalent.
- (1)  $A$  is completely reducible
  - (2)  $A$  has no proper large sub-module.
  - (3)  $L(A)$  is complemented.
- 
- b) If  $e^2 = e \in R$  and  $f^2 = f \in R$ , then  $eR \cong fR$  if and only if there exist  $u, v \in R$  such that  $vu = e$  and  $uv = f$ .
- Q4)** a) Prove that every free module is projective.
- b) Prove that,  $M$  is projective if and only if every epimorphism  $\pi : B \rightarrow M$  is direct.
- Q5)**
- a) If  $M$  is the direct product of a family of modules  $\{M_i / i \in I\}$ , then prove that,  $M$  is injective if and only if each  $M_i$  is injective.
- b) Prove that,  $M$  is injective if and only if  $M$  has no proper essential extension.